Section I: Prevalence of Smoking 2001, 2003, and 2005 Combined

	Never	Former	Curren	t	
Total	3	34	27	39	
Men Women		33 34	30 25	37 41	p < .01
vvoinen	`	D 4	25	41	
18-24	4	14	10	46	p < .0001
24-34	4	43	16	41	
35-44	3	35	19	46	
45-54	3	31	30	39	
55-64	2	26	37	37	
65+	3	30	48	22	
High cabaal ay laas	,	20	00	44	
High school or less		33	26	41	ns
More than high school	•	35	28	37	
Below median income	3	30	28	42	p < .0001
Above median income	4	40	30	30	•
Employed	3	37	26	37	p < .0001
Unemployed	2	28	20	53	•
Not in workforce	3	31	31	38	

Section II: Prevalence Chewing Tobacco Use 2001, 2003, and 2005 Combined

Men Only

	Never For	mer Cur	rent	
Men	57	30	13	
18-24 24-34 35-44 45-54 55-64	57 44 42 61 69 78	27 33 36 33 22 21	16 23 22 6 9 1	p < .0001
High school or less More than high school Below median income	59 55 58	29 31 29	12 14 13	ns
Above median income	53	29 34	13	ns
Employed Unemployed Not in workforce	59 54 68	28 28 24	14 18 8	p < .0001

Section III: Efforts and Intentions to Quit Smoking

2001, 2003, and 2005 Combined

	Tried to quit past year	Consider quitting next six months
Total	55	65
Men	46 p < .0001	61 ns
Women	61	67
18-24	76 p < .001	74 p < .05
24-34	65	65
35-44	58	69
45-54	44	62
55-64	47	59
65+	37	48
High school or less	56 ns	64 ns
More than high school	54	66
Below median income	57 ns	65 ns
Above median income	51	69
Employed	53 ns	66 ns
Unemployed	59	63
Not in workforce	56	64

Health care provider support for cigarette cessation

Advised to stop smoking 45 (2003 and 2005)

	2001	2003	2005
Recommended or offered NRT		45	53
Suggested setting a quit date			23
Sugested classes or counseling			20
Offered self help materials			35
Suggested aids other than NRT		34	

Strategies of Successful and Unsuccessful Quit Attempts in 2001

	Successful	Unsuccessful
NRT or prescription	8	21
Classes or counseling	3	7
Self-help materials	10	14
Telephone Quit Line	0	1

Section IV: Efforts and Intentions to Quit Using Chewing Tobacco

2001, 2003, and 2005 Combined

Men Only

	Tried to quit past year	Consider quiting in next six months
Men	47	64
18-24 24-34 35-44 45-54 55-64 65+	47 ns 52 50 36 36	84 ns 58 63 62 60
High school or less	37 p < .05	55 p < .05
More than high school	55	73
Below median income	49 ns	65 ns
Above median income	41	57
Employed	47 ns	59 ns
Unemployed	50	73
Not in workforce	45	77

Health care provider support for chewing tobacco cessation, 2001, 2003, 2005 combined

Advised to stop smoking 22 Recommended or offered NRT 22

Section V: Public Smoking Policies

	Smoking should be allowed Not at all Some area All areas				
Restaurants Bars, taverns, casinos Other indoor public areas	70 35 73	;	26 26 23	4 39 4	
	Smoking s Never smoked	should no Former smoker	t be permiti Current smoker	t	
Restaurants Bars, taverns, casinos Other indoor public areas	85 41 82		76 67 81	53 33 61	p < .0001
	In workpla		05 king should	d be allowed	
	Not at all	Some a	rea All area	19	
	85		12	3	
	Workplace	Policies	for Commo	on Areas	
	2001		20	005	ns
Not permitted at all	81			86	
Permitted in some areas	11			7	
Permitted in all areas	3			2	
No policy	6	•		4	
	Workplace	Policies	for Work A	reas	
	2001		20	005	ns
Not permitted at all	84	Ļ		85	
Permitted in some areas	8			7	
Permitted in all areas	3	3		4	
No policy	5	,		4	
	Exposed to second hand smoke in your work area in past week, 2005				
	Yes		No		
	11			89	

Section VI: Rules About Smoking in the Home

	Not at all	Some places	No restriction	
Total	59	16	25	
Men Women	58 60		28 22	ns
18-24 25-34 35-44 45-54 55-64 65+	63 70 58 57 56 57	15 21 16 11	15 21 26 33	p < .01
High school or less More than high school	53 66	_		p < .0001
Below median income Above median income	53 70	_		p < .0001
Children in home No children in home	66 50			p < .0001
Never smoked Former smoker Current smoker	78 72 38	11	15 17 37	p < .0001

Section VII: Second Hand Smoke

2003 and 2005

Second hand smoke is harmful or very harmful to one's health	92
Second hand smoke casues lung cancer in adults	77
Second hand smoke causes heart disease in adults	66
Second hand smoke causes respiratory problems in children	91
Second hand smoke causes Sudden Infant Death Syndrome	51
Exposed to second hand smoke in own home in past week	70
Exposed to second hand smoke in a car in past week	41